#### **Topic Area: Asthma**

#### AS-1: Reduce the death rate due to asthma

#### AS-1a: aged 35 to 64 years old

Target:	0.8 per 100,000 population
Baseline: (Year)	1.3 per 100,000 population (2005-2007)
Data source:	Death Certificate Database,
	Center for Health Statistics,
	New Jersey Department of Health

#### AS-1b: aged 65 years and older

Target:	2.0 per 100,000 population
Baseline: (Year)	3.7 per 100,000 population (2005-2007)
Data source:	Death Certificate Database,
	Center for Health Statistics,
	New Jersey Department of Health

### AS-2: Reduce the hospitalization rate due to asthma

#### AS-2a: among children aged under 5 years

Target:	328 per 100,000 children under age 5
Baseline: (Year)	469 per 100,000 children under age 5 (2009)
Data source:	Uniform Billing Patient Summary Data,
	Office of Health Care Quality Assessment,
	New Jersey Department of Health

#### AS-2b: among persons aged 5 to 64 years

Target:	112 per 100,000 persons aged 5-64 years (age-adjusted)
Baseline: (Year)	140 per 100,000 persons aged 5-64 years (age-adjusted) (2009)
Data source:	Uniform Billing Patient Summary Data,
	Office of Health Care Quality Assessment,
	New Jersey Department of Health

#### AS-2c: among persons aged 65 years and older

Target:	260 per 100,000 persons aged 65 years and older (age-adjusted)
Baseline: (Year)	325 per 100,000 persons aged 65 years and older (age-adjusted) (2009)
Data source:	Uniform Billing Patient Summary Data,

Office of Health Care Quality Assessment,
New Jersey Department of Health

#### AS-3: Reduce the rate of Emergency Department (ED) visits due to asthma

#### AS-3a: among children aged under 5 years

Target:	1220 per 100,000 children under age 5
Baseline: (Year)	1525 per 100,000 children under age 5 (2009)
Data source:	Uniform Billing Patient Summary Data,
	Office of Health Care Quality Assessment,
	New Jersey Department of Health

#### AS-3b: persons aged 5 to 64 years

Target:	501 per 100,000 persons aged 5-64 years (age-adjusted)
Baseline: (Year)	626 per 100,000 persons aged 5-64 years (age-adjusted) (2009)
Data source:	Uniform Billing Patient Summary Data,
	Office of Health Care Quality Assessment,
	New Jersey Department of Health

#### AS-3c: persons aged 65 years and older

Target:	128 per 100,000 persons aged 65 years and older (age-adjusted)
Baseline: (Year)	160 per 100,000 persons aged 65 years and older (age-adjusted)
	(2009)
Data source:	Uniform Billing Patient Summary Data,
	Office of Health Care Quality Assessment,
	New Jersey Department of Health

# AS-4: Reduce the proportion of persons with asthma who miss school or work days

## AS-4a: Children attending school or preschool with asthma who miss days because of asthma

Target:	To be determined
Baseline: (Year)	Not yet available (2013)
Data source:	New Jersey Behavioral Risk Factor Asthma Call-back Survey,
	Center for Health Statistics,
	New Jersey Department of Health

## AS-4b: Adults aged 18 years and older with asthma who miss work days or limit usual activities because of asthma

Target:	To be determined
Baseline: (Year)	Not yet available (2013)
Data source:	New Jersey Behavioral Risk Factor Asthma Call-back Survey,
	Center for Health Statistics,
	New Jersey Department of Health

# AS-5: Increase the proportions of persons with asthma who have ever received an asthma action plan or asthma management plan from a health professional

#### AS-5a: Children aged 0 to 17 years

Target:	To be determined
Baseline: (Year)	Not yet available (2013)
Data source:	New Jersey Behavioral Risk Factor Asthma Call-back Survey,
	Center for Health Statistics,
	New Jersey Department of Health

#### AS-5b: Adults aged 18 years and older

Target:	To be determined
Baseline: (Year)	Not yet available (2013)
Data source:	New Jersey Behavioral Risk Factor Asthma Call-back Survey, Center for Health Statistics, New Jersey Department of Health

AS-6: Increase the proportion of persons with asthma who have ever been advised by a health professional to change things in the home, school, or work to improve their asthma.

AS-6a: Children aged 0 to 17 years

Target:	To be determined
Baseline: (Year)	Not yet available (2013)
Data source:	New Jersey Behavioral Risk Factor Asthma Call-back Survey, Center for Health Statistics,
	New Jersey Department of Health

## AS-6b: Adults aged 18 years and older

Target:	To be determined
Baseline: (Year)	Not yet available (2013)
Data source:	New Jersey Behavioral Risk Factor Asthma Call-back Survey,
	Center for Health Statistics,
	New Jersey Department of Health